

---

# M & E: Partnerships and Lessons Learned

*Niamh Darcy, POPPHI M&E lead*

20 November 2009



**USAID**  
FROM THE AMERICAN PEOPLE



**POPPHI**

Prevention of Postpartum  
Hemorrhage Initiative

# Background

- Discussions Nov 2006 on ‘standardized’ USAID AMTSL indicator
- Development of Indicator Reference Sheet

## AMTSL INDICATOR REFERENCE SHEET – V11 Apr 5<sup>th</sup> 2007

**Indicator 1:** Number and Percentage of women in facilities and home where the woman received active management of the third stage of labor (AMTSL) by skilled birth attendants (SBAs)<sup>1</sup> within a specified time period

### DESCRIPTION

**Precise Definition:** Number and percent of women in facilities and homes where the woman received AMTSL by SBAs in targeted areas in a specified time period. This includes vaginal deliveries only.<sup>2</sup> Targeted areas are those where the United States Agency for International Development partner and Cooperating Agency (CA) maternal and child health projects are implementing AMTSL interventions – these include public and private health facilities, rural and urban health facilities, as well as home births with SBAs. AMTSL is defined as the following three elements:

- a. Use of uterotonic drug within one minute of birth (oxytocin is the drug of choice, preferred 10 IU/IM).
- b. Performance of controlled cord traction.
- c. Performance of uterine massage after the delivery of the placenta.



**USAID**  
FROM THE AMERICAN PEOPLE



**POPPHI**

Prevention of Postpartum  
Hemorrhage Initiative

# AMTSL M&E

- AMTSL national indicator – regular AMTSL data collection and use
- AMTSL advocacy at district, regional and national level HMIS staff
- ATMSL M&E capacity building



# AMTSL M&E Capacity Building

- Emphasize importance of AMTSL data collection as part of AMTSL training – reinforce practice
- Emphasize importance of AMTSL supportive supervision
- Emphasize importance of using the AMTSL data within the facility to improve AMTSL practice



**USAID**  
FROM THE AMERICAN PEOPLE



**POPPHI**

Prevention of Postpartum  
Hemorrhage Initiative

# Recording AMTSL practice in Facilities (1)

- AMTSL data recorded in
  - Partograph
  - patient record
  - delivery register/log
- All three components separately or single marker
- Use of stamps, pre-printed records, hand written

# Recording AMTSL practice in Facilities (2)

- Mali, Ghana, DRC, Benin: Partograph (pre-printed) and Delivery log
- Pakistan: Partograph and Delivery log
- Bangladesh: Patient's chart (stamp) and Delivery log (stamp/hand written)
- Indonesia, Nicaragua: Partograph (pre-printed, 3 components)

# Recording AMTSL practice in Facilities (3)

Duration of Stage III:..... minutes

Oxytocin 10 U IM

Yes, time: ... min. after delivery

No, explain...

2nd dose of oxytocin?

Yes, explain.....

No

Controlled cord traction

Yes


No, explain.....

Fundal Massage:

Yes


No, explain.....

AMTSL	
10 unit Oxytocin provided	<input type="checkbox"/>
Controlled Cord Traction	<input type="checkbox"/>
Uterine massage	<input type="checkbox"/>



# PPH Incidence

- Diagramming AMTSL and PPH incidence in facilities



# Supportive Supervision

- MoH AMTSL integrated into Supportive Supervision
  - Pakistan, Ukraine, Bangladesh: Project based
  - Mali: Project based, but planned with MoH for 2010
  - Nicaragua (MoH)
  - Senegal (added to national RH supervision tools)
  - Mali, Benin, DRC: Proposed to integrate into RH SS tools once AMTSL indicator adopted

# Recording AMTSL in HMIS (district, regional, national)

---

- Plans: Mali & DRC (Plans for 2010), Benin [national HMIS]
- Ukraine, Bangladesh (project based)
- Senegal, Nicaragua, Honduras [national HMIS]

