

Additional References

AMTSL

- WHO recommendations (Available at: http://www.who.int/making_pregnancy_safer/publications/WHORecommendationsforPPHaemorrhage.pdf)::
- Bagley C. A comparison of 'active' and 'physiological' management of the third stage of labour. *Midwifery* 1990; 6: 3-17.
- Dombrowski MP, Bottoms SF, Saleh AA, Hurd WW, Romero R. Third stage of labor: analysis of duration and clinical practice. *American Journal of Obstetrics and Gynecology*. 1995;172:1279–84.
- Everett F, Magann EF, Evans S, Chauhan SP, Lanneau G, Fisk AD, Morrison JC. The Length of the Third Stage of Labor and the Risk of Postpartum Hemorrhage. *Obstetrics & Gynecology*. 2005;105(2): 290–293.
- JHPIEGO. Preventing Postpartum Hemorrhage: Active Management of the Third Stage of Labor—A Maternal And Neonatal Health Program Best Practice. *JHPIEGO TrainerNews*. Washington, DC: JHPIEGO; November 2001. Available at: <http://www.reproline.jhu.edu/english/6read/6issues/6jtn/v4/tn110hemor.htm>.
- Stephenson P. Active Management of the Third Stage of Labor: A Simple Practice to Prevent Postpartum Hemorrhage. USAID Global Health Technical Brief. June 2005. MAQ website. Available at: <http://www.maqweb.org/techbriefs/tb13activemgmt.shtml>.

AMTSL practiced at the community / home level

- Sanghvi H, Wiknjosastro G, Chanpong G, Fishel J, Ahmed S, Zulkarnain M. *prevention of postpartum hemorrhage study West Java, Indonesia*. Baltimore, MD: JHPIEGO. 2004. (Available at: <http://www.jhpiego.org/resources/pubs/mnh/pphjavastudy.pdf>)
- Strand R, Da Silva F, Jangsten E, Bergström S. Postpartum hemorrhage: a prospective, comparative study in Angola using a new disposable device for oxytocin administration. *Aca Obstetricia et Gynecologica Scandinavica*. 2005;84:260–265
- Tsu VD, Mai TTP, Nguyen HD, and Luu HTT. Reducing postpartum hemorrhage in Vietnam: Assessing the effectiveness of active management of third-stage labor. *J. Obstet. Gynaecol. Res*. October 2006; Vol. 32 (5): 489–496.
- Tsu VD, Sutanto A, Vaidya K, Coffey P, Widjaya A. Averting maternal death and disability: Oxytocin in prefilled Uniject™ injection devices for managing third-stage labor in Indonesia. *International Journal of Gynecology and Obstetrics*. 2003; 83: 103–111
- Walraven G, et al. Misoprostol in the management of the third stage of labour in the home delivery setting in rural Gambia: a randomized controlled trial. *BJOG: an International Journal of Obstetrics and Gynaecology*. 2005;112:1277–1283.

Choice of uterotonic

- WHO recommendations (Available at: http://www.who.int/making_pregnancy_safer/publications/WHORecommendationsforPPHaemorrhage.pdf)
- FIGO/ICM recommendations (Available at: http://pphprevention.org/files/FIGO-ICM_Statement_November2006_Final.pdf):
- Bamigboye A et al. Randomized comparison of rectal misoprostol with syntometrine for management of third stage of labor. *Acta Obstet Gynecol Scand* 1998; 77: 178–181.
- Cotter A, Ness A, Tolosa J (12) Prophylactic oxytocin for the third stage of labour. *Cochrane Database Syst. Rev.* 2001, Issue 4.
- El-Refaey H et al. The misoprostol third stage of labour study: A randomized controlled comparison trial between orally administered misoprostol and standard management. 2000; 107: 1104-1110.
- Hogerzeil HV, Walker GJ (4) Instability of (methyl)ergometrine in tropical climates: an overview. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 1996;69:25-9.
- McDonald et al. Randomized controlled trial of oxytocin alone versus oxytocin and ergometrine in active management of third stage of labor. *BMJ* 1993; 307(6913):1167–1171.
- Ng PS et al. A multicentred randomized controlled trial of oral misoprostol and IM syntometrine in the management of third stage of labour. *Hum Reprod.* 2001; 16: 31-35.
- Okan Ozkaya O, Sezik M, Kaya H, Desdicioglu R, Dittrich R. Placebo-controlled randomized comparison of vaginal with rectal misoprostol in the prevention of postpartum hemorrhage. *Journal of Obstetrics and Gynaecology Research.* 2005; 31 (5): 389-393. (Available at:<http://www.blackwell-synergy.com/doi/abs/10.1111/j.1447-0756.2005.00307.x>)
- Sorbe B (13) Active pharmacologic management of the third stage of labor. A comparison of oxytocin and ergometrine. *Obstet.Gynecol.* 1978;52:694-7.

Misoprostol for prevention of PPH

- FIGO/ICM recommendations (Available at: http://pphprevention.org/files/FIGO-ICM_Statement_November2006_Final.pdf):
- Broekhuizen FF. Misoprostol, its use in obstetrics and gynaecology. Current problems in Obstetrics, Gynaecology, and Fertility. 2000; 23: 183-224.
- Carpenter JP. Misoprostol for Prevention of Postpartum Hemorrhage: An Evidence-Based Review by the US Pharmacopeia, Rockville, Maryland: United States Pharmacopeia, 2001.
- Derman R, et al. Oral misoprostol in preventing postpartum haemorrhage in resource-poor communities: a randomized controlled trial. *The Lancet*. 2006;368:1248–1253.
- El-Refaey H et al. Use of misoprostol in prevention of postpartum hemorrhage. *British Journal of Obstetrics and Gynaecology*. 1997; 104: 336-339.
- El-Refaey H et al. The misoprostol third stage of labour study: A randomized controlled comparison trial between orally administered misoprostol and standard management. 2000; 107: 1104-1110.
- Goldberg AB, Greenberg MA, Darney PD. Misoprostol and pregnancy. *New England Journal of Medicine* 2001; 344:38-47.
- Gulmezoglu, AM et al. (15) Prostaglandins for prevention of postpartum haemorrhage. *Cochrane Database. Syst.Rev* 2006, Issue 4.
- Høj L, Cardoso P, Nielsen BB, et al. Effect of sublingual misoprostol on severe postpartum haemorrhage in a primary health centre in Guinea-Bissau: randomised double blind clinical trial. *British Medical Journal*. 2005;331:723.
- Jadesimi A, Okonofua FE. Tackling the unacceptable: Nigeria approves misoprostol for postpartum haemorrhage. *J Fam Plann Reprod Health Care*. 2006;32:213-4
- Langenbach C. Misoprostol in preventing postpartum hemorrhage: A meta-analysis. *International Journal of Gynecology & Obstetrics*. 2006;92:10–18.
- Lumbiganon P et al. Misoprostol dose-related shivering and pyrexia in third stage of labour. Who collaborative trial of misoprostol in management of the third stage of labour. *British Journal of Obstetrics and Gynaecology*. 1999; 106: 304-308.
- Ng PS et al. A multicentred randomized controlled trial of oral misoprostol and IM syntometrine in the management of third stage of labour. *Hum Reprod*. 2001; 16: 31-35.
- Prata, N. A global overview of ongoing misoprostol studies. *International Journal of Gynecology and Obstetrics* (2006), 94 (Supplement 2), S147-S148.
- Sanghvi H, Wiknjosastro G, Chanpong G, Fishel J, Ahmed S, Zulkarnain M. *prevention of postpartum hemorrhage study West Java, Indonesia*. Baltimore, MD: JHPIEGO. 2004. (Available at: <http://www.jhpiego.org/resources/pubs/mnh/pphjavastudy.pdf>)
- Walraven G, et al. Misoprostol in the management of the third stage of labour in the home delivery setting in rural Gambia: a randomized controlled trial. *BJOG: an International Journal of Obstetrics and Gynaecology*. 2005;112:1277–1283.

Uterotonics without CCT

- WHO recommendations (Available at: http://www.who.int/making_pregnancy_safer/publications/WHORecommendationsforPPHaemorrhage.pdf)
- Derman R, et al. Oral misoprostol in preventing postpartum haemorrhage in resource-poor communities: a randomized controlled trial. *The Lancet*. 2006;368:1248–1253.
- Jackson KW, Allbert JR, Schemmer GK, Elliot M, Humphrey Angela MS; Taylor, Jane RN. A randomized controlled trial comparing oxytocin administration before and after placental delivery in the prevention of postpartum hemorrhage. *American Journal of Obstetrics & Gynecology*. October 2001;185(4):873-877. (Available at: http://www.medscape.com/medline/abstract/11641669?src=emed_ckb_ref_0)
- Prata, N. A global overview of ongoing misoprostol studies. *International Journal of Gynecology and Obstetrics* (2006), 94 (Supplement 2), S147-S148.
- Sanghvi H, Wiknjosastro G, Chanpong G, Fishel J, Ahmed S, Zulkarnain M. *prevention of postpartum hemorrhage study West Java, Indonesia*. Baltimore, MD: JHPIEGO. 2004. (Available at: <http://www.jhpiego.org/resources/pubs/mnh/pphjavastudy.pdf>)
- Walraven G, et al. Misoprostol in the management of the third stage of labour in the home delivery setting in rural Gambia: a randomized controlled trial. *BJOG: an International Journal of Obstetrics and Gynaecology*. 2005;112:1277–1283.

Delayed cord clamping

- WHO recommendations: Available at: http://www.who.int/making_pregnancy_safer/publications/WHORecommendationsforPPHaemorrhage.pdf)
- Ceriani Cernadas JM et al. The effect of timing of cord clamping on neonatal venous hematocrit values and clinical outcome at term: a randomized, controlled trial. *Pediatrics*. 2006;117:e779-e786.
- Chaparro CM et al. Effect of timing of umbilical cord clamping on iron status in Mexican infants: a randomised controlled trial. *Lancet* 2006;367:1997-2004.
- Emhamed MO, van Rheenen P, Brabin BJ. The early effects of delayed cord clamping in term infants born to Libyan mothers. *Trop.Doct.* 2004;34:218-22.
- Gupta R, Ramji S. Effect of delayed cord clamping on iron stores in infants born to anemic mothers: a randomized controlled trial. *Indian Pediatrics*. 2002;39(2):130–135.
- Mercer JS et al. Delayed cord clamping in very preterm infants reduces the incidence of intraventricular hemorrhage and late-onset sepsis: a randomized, controlled trial. *Pediatrics* 2006;117:1235-42.
- Rabe H, Reynolds G, Diaz-Rossello J. Early versus delayed umbilical cord clamping in preterm infants. *Cochrane Database Syst.Rev.* 2004; Issue 4.
- Van Rheenen P, Brabin BJ. Late umbilical cord-clamping as an intervention for reducing iron deficiency anaemia in term infants in developing and industrialized countries: a systematic review. *Ann.Trop.Paediatr.* 2004;24:3-16.
- Yao AC, Moinian M, Lind J. Distribution of blood between infant and placenta after birth. *Lancet*. 1969;7626:871–873

Nipple stimulation to increase uterine contractions during the third stage of labour

- Irons DW, Sriskandabalan P, Bullough CHW. A simple alternative to parenteral oxytocics for the third stage of labor. *Int J Obstet Gynecol* 1994; 46:15–18.

CCT without uterotonics

- Not recommended.

Uterine massage

- Abdel-Aleem H, Hofmeyr G, Shokry M, El-Sonoosy E. Uterine massage and postpartum blood loss. *International Journal of Gynecology & Obstetrics*. 93 (3): 238-239
- Hofmeyr GJ, Abdel-Aleem H. Uterine massage for preventing postpartum haemorrhage. (Protocol) *Cochrane Database of Systematic Reviews* 2007, Issue 2. Art. No.: CD006431. DOI: 10.1002/14651858.CD006431.

Use of the risk approach

- De Groot ANJA, Slort W, Van Roosmalen J. Assessment of the risk approach to maternity care in a district hospital in rural Tanzania. *Int J Gynaecol Obstet* 1993; 40:33-37.
- Fortney J. Antenatal risk screening and scoring: a new look. *Int J Gynecol Obstet* 1995; 50(Suppl 2): S53–S58.
- Yuster EA. Rethinking the role of the risk approach and antenatal care in maternal mortality reduction. *Int J Gynecol Obstet* 1995; 50(Suppl 2): S59–S61.